

## Sheet 6

### Exercise 1.1 (Weak and strong convergence)

Let  $\mathcal{H}$  be a Hilbert space and  $(f_n)_{n \in \mathbb{N}}$  a sequence of  $\mathcal{H}$  that converges weakly to  $f$  in  $\mathcal{H}$  and such that  $(\|f_n\|_{\mathcal{H}})_{n \in \mathbb{N}}$  converges to  $\|f\|_{\mathcal{H}}$ . Show that  $(f_n)_{n \in \mathbb{N}}$  converges strongly to  $f$  in  $\mathcal{H}$ .

### Exercise 1.2 (Density in $H^s(\mathbb{R}^d)$ )

Let  $s \in \mathbb{R}$ . Show that  $\mathcal{S}(\mathbb{R}^d)$  is dense in  $H^s(\mathbb{R}^d)$ .

### Exercise 1.3 (Local compact embedding)

Let  $t < s$ ,  $\varphi \in \mathcal{S}(\mathbb{R}^d)$ . The goal of this exercise is to show that the multiplication by  $\varphi$  is a compact operator from  $H^s(\mathbb{R}^d)$  to  $H^t(\mathbb{R}^d)$ .

Let  $(u_n)_{n \in \mathbb{N}}$  be a sequence of  $H^s(\mathbb{R}^d)$  such that  $\sup_{n \in \mathbb{N}} \|u_n\|_{H^s} \leq 1$ .

1. Show that, up to extraction of a subsequence,  $(u_n)_{n \in \mathbb{N}}$  converging weakly in  $H^s(\mathbb{R}^d)$  to an element  $u$ .

Let us set  $v_n := u_n - u$ .

2. Show that there exists a constant  $C_1$  such that

$$\sup_{n \in \mathbb{N}} \{\|\varphi v_n\|_{H^s}\} \leq C_1.$$

3. Show that for any positive real number  $R$ , we have

$$\|\varphi v_n\|_{H^t} \leq \int_{B(0,R)} (1 + |\xi|^2)^t |\mathcal{F}(\varphi v_n)|^2 d\xi + \frac{C_1^2}{(1 + R^2)^{s-t}}.$$

Let us consider  $\varepsilon > 0$ .

4. Show that there exists a positive real number  $R$  such that

$$\frac{C_1^2}{(1 + R^2)^{s-t}} \leq \varepsilon.$$

5. For all  $\xi \in \mathbb{R}^d$ , we set  $\psi_\xi := \mathcal{F}^{-1}((1 + |\cdot|^2)^{-s} \mathcal{F}(\varphi)(\xi - \cdot))$ . Show that, for any  $\xi \in \mathbb{R}^d$ ,  $\psi_\xi$  belongs to  $\mathcal{S}(\mathbb{R}^d)$  and that

$$\forall \xi \in \mathbb{R}^d, \quad \mathcal{F}(\varphi v_n)(\xi) = \langle \psi_\xi, v_n \rangle.$$

6. Deduce that for any  $\xi \in \mathbb{R}^d$ , we have  $\lim_{n \rightarrow +\infty} \mathcal{F}(\varphi v_n)(\xi) = 0$ .

7. Assume that there exists a positive real number  $M > 0$  such that

$$\sup_{\xi \in B(0,R), n \in \mathbb{N}} \{|\mathcal{F}(\varphi v_n)|\} \leq M. \quad (1)$$

8. Conclude.

We will now show (1).

9. Show that there exists a positive real number  $C_2$  such that

$$\forall \mu \in \mathbb{R}^d, \quad |\widehat{\varphi}(\mu)| \leq \frac{C_2}{(1 + |\mu|^2)^{\frac{d}{2} + |s| - 1}}.$$

10. Show that for any  $\xi \in B(0, R)$ ,

$$\begin{aligned} & \int_{\mathbb{R}^d} (1 + |\eta|^2)^{-s} |\widehat{\varphi}(\xi - \eta)|^2 d\eta \\ & \leq C_1 \int_{|\eta| \leq 2R} (1 + |\eta|^2)^{|s|} ds + C_2 \int_{|\eta| \geq 2R} \frac{(1 + |\eta|^2)^{|s|}}{(1 + |\xi - \eta|^2)^{\frac{d}{2} + |s| + 1}} d\eta. \end{aligned}$$

11. Deduce that there exists a positive real number  $C_3$  such that

$$\forall \xi \in B(0, R), \quad \int_{\mathbb{R}^d} (1 + |\eta|^2)^{-s} |\widehat{\varphi}(\xi - \eta)|^2 d\eta \leq C_3 (1 + R^2)^{|s| + \frac{d}{2}}$$

(*Hint*: to bound  $\int_{|\eta| \geq 2R} (1 + |\eta|^2)^{-s} (1 + |\xi - \eta|^2)^{-(\frac{d}{2} + |s| + 1)} d\eta$ , use that if  $|\xi| \leq R$  and  $|\eta| \geq 2R$ , we have  $|\xi - \eta| \geq \frac{|\eta|}{2}$ .)

12. Deduce that (1) holds.

#### Exercise 1.4 (Norm of the heat propagator)

Let  $t > 0$ . Show that  $\|e^{t\Delta}\|_{\mathcal{B}(L^2(\mathbb{R}^d))} = 1$ .